

# Journal of Community Medicine and Health Solutions

Volume - 3, Issue - 1

**Opinion**      **Published Date:-2022-12-27 16:36:17**

[The role of health sectors in the fighting misconception of COVID-19 vaccine](#)

In the long run people around the globe particularly those who live in underdeveloped Countries in Africa, especially in Ethiopia a large number of communities has a misconception that disease may be transmitted by utilizing health service due to a lack of trust in health service delivery system, Most of the population strongly related with religion, they claimed that it is sin or God or Allah driven. According to a health behavior model, knowledge and practice are contributing factors for increasing healthcare seeking among patients with low levels of trust [1,2].

---

**Short Review**      **Published Date:-2022-12-01 12:32:51**

[Association Between Internet Gaming Disorder And Attention Deficit Hyperactivity Disorder: A Narrative Review](#)

Background: The rapid increase in the popularity of internet use and video games is associated with multiple consequences and comorbidities; therefore, investigating this issue is important, especially now that the number of people playing video games has reached approximately 2.2 billion people worldwide.  
Method: This paper reviews the existing literature on Google Scholar and PubMed to examine the association between internet gaming disorder (IGD) and attention deficit hyperactivity disorder (ADHD) between 2013 and 2020.  
Result: The reviewed studies reveal a statistically significant association between ADHD and IGD. The reviewed studies demonstrate that ADHD is a predictive factor for the development and persistence of IGD and affects the severity of IGD symptoms.  
Conclusion: Especially among those with ADHD, IGD causes life distress and social life impairment. This review indicates a statistically significant association between ADHD and IGD; however, longitudinal studies are needed to assist the direction of this association and make relevant and appropriate recommendations to prevent its consequences and comorbidities.

---

**Research Article**      **Published Date:-2022-10-13 12:49:11**

[Experience in optimizing the accessibility of services for tuberculosis in the Republic of Tajikistan](#)

Many researchers in their publications on tuberculosis (TB) in high-burden countries recommended the integrated work of different ministries and departments with the involvement of non-governmental public associations, volunteers, and activists, including former patients [1-8].

---

**Opinion**      **Published Date:-2022-09-02 16:42:14**

[Effect of COVID-19 on the ambient air pollution in Tunisia during 2020 and 2021 years](#)

The epidemic of COVID-19 was reported in Wuhan, China in December 2019 and turned into a national crisis, with infected individuals diagnosed all over China [1-3]. In early March 2020, the World Health Organization (WHO) declared that the Wuhan epidemic has turned into a global pandemic. Many European countries have started to know several cases affected by this coronavirus, which is known to be highly contagious. The WHO has launched several recommendations to curb the spread of this virus and to call the general confinement establishment in the affected countries.

---

[The factors associated with mental health and the impact of COVID-19](#)

The mental health impact caused by COVID-19 on adolescents was reviewed, and due to limited data, adult results were included, to support our assertion that additional mental health resources are needed for both adult and young people. Positive gains would include improved socio-emotional skills, a decrease in maladaptive behaviors contributing to the disruption in interpersonal relationships and lifetime achievements, suicide attempts and psychopathology, persistent mental health concerns found in the juvenile justice and foster care systems and substance use addiction later in adulthood (The American Psychological Association, 2019; Garber & Weersing, 2010; Office of Juvenile Justice and Delinquency Prevention, n.d.; Tomasello, 2018;). Geiger & Davis, 2019, found that 13% or 3.2 million United States (U.S.) teenagers aged 12 - 17 years old experienced at least one major depressive episode with the depression rate increasing 59% from 2007 to 2017. Therefore, we assert that developing programs to overcome barriers to mental health aid can reduce instances experienced in adolescence and adulthood.

---

[Face mask utilization and associated factors in combating COVID-19 pandemic among government employee in Akaki district administration offices, Oromia, Ethiopia, 2022](#)

Background: The novel coronavirus is a rapidly spreading respiratory disease that has been declared a pandemic by the World Health Organization (WHO) and a global public health emergency. The use of face masks has been recommended by the WHO and the Centers for Disease Control (CDC) as a standard prevention method for transmission of COVID-19.

Objective: The objective of this study is to determine face mask utilization and associated factors in combating the COVID-19 pandemic among government employees in Akaki district administration offices in Akaki District, Oromia, Ethiopia, 2022

Methods: A quantitative cross-sectional study was conducted from December 1, 2021, to February 15, 2022, on 385 government employees working in Akaki district administration offices. After obtaining consent from the study participants, data were collected using pretested, self-administered, and standardized questionnaires adapted from other studies. After the data was collected, it was entered into Epi info version 7.2.6, cleaned, and analyzed using SPSS version 26. A logistic regression model was computed to measure the association between the predictor and outcome variables. A p - value of .05 with a 95% CI was used as the cut-off point to declare the level of statistical significance.

Results: The study showed that the magnitude of good practice for facemask utilization was 213 (53.5%) (95% CI: 1.50, 1.60) for preventing COVID-19. In the multivariate logistic regression analysis, the odds of using face masks among male employees (AOR = 0.275; 95% CI: 0.137, 0.555), employees aged 20-29 (AOR = 0.17; 95% CI: 0.065, 0.481 ), employees aged 30-39 (AOR = 0.260, 95% CI: 0.109, 0.623), employees of less than five family size (AOR = 0.549, 95% CI: 0.303, 0.995), work experience at 6-12 (AOR = 0.32, 95 CI: 0.120, 0.450), poor knowledge about face mask use (AOR = 0.504, 95% CI: 0.302, 0.844), and employees with a negative attitude (AOR = 0.430, 95% CI: 0.256, 0.721) were factors significantly associated with face mask utilization.

Conclusion: The magnitude of facemask utilization was low (53.5%) compared to other studies. The results of the study showed that age, sex, household family size, work experience, poor knowledge, and the negative attitude of employees toward facemask utilization were among the factors significantly associated with facemask utilization.

---

[Associations of Exercise Habits and Circulatory Dynamics with Peripheral Lower Limb Body Composition in Healthy Community-dwelling Older Individuals](#)

We investigated the associations between circulatory dynamics and Body Composition (BC) of the peripheral lower limbs with Exercise Habits (EHs). We surveyed the EHs of 34 healthy older adults (6 men, 28 women) and measured their BC and skin perfusion (SP). SP did not correlate significantly with any other variable. Participants with established EHs had a lower body weight and Fat Tissue thickness (Fat-T) than those without. Multiple logistic regressions with the presence or absence of an EH as the dependent variable and BC values as the independent variables established Fat-T as a significant variable. Sex was independently associated with soleus muscle thickness, Fat-T, Bone Density (BD), and leg segmental water. Age was independently associated with Fat-T and BD, and body mass index was associated with Fat-T. EHs were not independently associated with any of the BC measurements. Physical characteristics and EHs have minimal influence on peripheral circulatory dynamics.

---

[The effects of religiosity and knowledge on the perceptions and attitude of muslim doctors that influence the intentions that impact on behavior in using halal drugs in Indonesia](#)

Introduction: In the spirit of maintaining health and seeking medicine to cure a disease, the question of halal and haram still needs to be answered. This coincides with the words of the Prophet Muhammad SAW "Verily Allah does not make your healing in what He has forbidden you" (HR. Imam Bukhari). The purpose of this quantitative study was to determine the effect of religiosity, perception, knowledge, and attitudes of Muslim doctors on the use of halal drugs in Indonesia.

Materials and methods: This quantitative research method with correlational type examines the relationship between religiosity, perceptions, knowledge, attitudes, intentions, and behaviors that are shared online using google form. Data was measured using a Likert scale. A five-point Likert scale was used for statements: Strongly Disagree: 1, Disagree: 2, Uncertain: 3, Agree: 4, and Strongly Agree: 5. The sample size used was 435 Muslim doctors. The data obtained were then processed using the SmartPLS 3.0 application.

Results: The results obtained are that there is a relationship between the religiosity of Muslim doctors on perceptions so as to encourage intention so that there is a change in behavior in the use of halal drugs in Indonesia. In addition, there is a relationship between knowledge and attitudes so as to encourage intentions that cause behavioral changes in the use of halal drugs in Indonesia by Muslim doctors. In fact, every practice depends on the intention, therefore it is important to maintain the intention, which in this study is influenced by perceptions and attitudes so that charity will result in commendable behavior that is blessed by Allah SWT. In addition, there is also a need for education about halal drugs that are appropriate to be taught at the medical faculty of Islamic universities in Indonesia.

---

[Indoor air pollution and behavioral factors affecting to COVID-19 transition](#)

Environmental and behavioral factors are very important for exposure to airborne SARS-CoV-2. Indoor environments are related to infection events, including super-spreader events and outbreaks. Indoor, poorly ventilated, and crowded areas, such as restaurants, cinemas, and bars can be effective in the accumulation of aerosols full of viruses, especially if people are in conversations and stay there for a long time period. At longer distances (more than 1.5 meters), small aerosols that can stay in the air for a longer period of time are dominant. The super-spreader events in which people have been infected at a distance away show that this remote transmission occurs. The exposure risk to longer intervals is likely to be more in domestic environments and indoor spaces that lack sufficient ventilation. Layer interventions are of fundamental importance. Therefore, it is important to take preventive measures as much as possible and follow them as carefully as possible, because no intervention alone will be effective in eliminating the risk. These include spacing, lining, hand hygiene, filtration, and ventilation.

---

[Risk of activation of human viruses lurking in ambient following COVID-19 prevention supplies excessive use](#)

Due to extensive COVID-19 prevention measures, millions of tons of chemicals penetrated the natural environment. Alterations of viruses in the environment, the neglected perceiver of environmental fluctuations, remain obscure. Chemicals especially trihalomethane restrained the virus community diversity. Segments of SARS-CoV-2 RNA have been detected near hospitals that suggesting the environment as a missing link in the transmission route. Human viruses lurking in the environment were potentially activated by pandemic prevention chemicals, warning an overlooked burden to human health. This letter warns of the risk of activation of human viruses in the environment following the overuse of COVID-19 prevention devices and emphasizes the long-term monitoring of environmental viruses in the post-pandemic period.

---

[Knowledge, attitude and practice towards facemask use amid the COVID-19 pandemic among public transport drivers in Debre-Tabor town, Ethiopia](#)

**Background:** For the prevention of respiratory infection similar to the current pandemic COVID-19, face masks are considered an effective method. Coronavirus disease (COVID-19) is a highly transmittable virus that continues to disrupt livelihoods, particularly those of low-income segments of society, around the world has poor knowledge towards facemask at the same time has huge gape towards attitude which affects the behaviors. Therefore this study was aimed to assess knowledge, attitude, and practice towards facemask use amid the COVID-19 pandemic among public transport drivers in Debre-Tabor town, Ethiopia, 2021.

**Methods:** A cross-sectional study was conducted among 407 public transport drivers in Debre-Tabor town from July to August 2021. The study participants were selected using a simple random sampling technique after proportionally allocating the sample size from the total number of public transport drivers, and finally, study subjects were elected continuously working in Debre- Tabor Town public transport drivers. The data were collected by trained data collectors using a structured questionnaire with face-to-face interview administered questioners and an on-the-spot observational checklist. The collected data were checked, coded, and entered into SPSS version 23, and for data cleaning and analysis.

**Result:** A total of 407 drivers responded to the questionnaire with a response rate of 96.4% were participated in this study. The mean age was 32.75 years (range: 19-45 years). Among them majority was male 383(94.1%) and the majority of 177(43.5%) were Bajaj drives. Among the responders, 114(28%) used radio to gather information, and 41(10.1%) drivers where were caught by COVID-19, but, only 30(7.4%) were vaccinated COVID-19 vaccine.

Knowledge, attitude and practices regarding facemask use among these drivers were 368(90.4% [95% CI: 87.7-93.1]), 163(40% [95% CI: 37.5% - 43.2%]), and 23 ((5.6%) [95% CI: 3.7% - 8.3%]) Regarding facemask use against COVID-19 infection respectively

**Conclusion and recommendation:** The current study was able to demonstrate 90.4% of drivers had good knowledge. Positive attitude and practice levels of face mask use among drivers were less than half of them or 40%, and 5.6% respectively. There is a high supply of Surgical Face masks but poor attitude and practice. Almost all of the participants don't wear a face mask despite having good knowledge about the transmission routes of COVID-19. The government COVID-19 prevention task force or health sector should be continued their awareness creation to bring behavioral change for the implementation of the COVID-19 prevention method, especially facemask.

---